

PHYTOCHEMICAL AND ANTIMICROBIAL EVALUATIONS OF SOME MEDICINAL PLANTS IN SURIGAO DEL SUR, PHILIPPINES

Gemma A. Gruyal

¹ Surigao del Sur State University-Cantilan Campus
Department of Biological Science
College of Teacher Education
Cantilan, Surigao del Sur 8317, Philippines

*For correspondence; Tel. 09088838950, E-mail: gemma_gruyal@yahoo.com

ABSTRACT: Medicinal plants have been widely used since before recorded history for curing and preventing various ailments. The bioactive chemical constituents and antimicrobial activity of five medicinal plants against *S. aureus* and *E. coli* were evaluated. A qualitative phytochemical analysis was performed for the detection of carbohydrates, reducing sugars, saponins, flavonoids, steroids, alkaloids and glycosides. Results revealed that *M. arvensis* and *C. frutescens* were found to have greater number of secondary metabolites. The ethanolic extract of *M. arvensis*, *C. frutescens* and *C. aromaticus* showed significant antimicrobial activity against *S. aureus*. Extracts of *C. citratus*, *C. frutescens* and *C. aromaticus* were also found to be effective against *E. coli*. The occurrence of these bioactive metabolites in the five medicinal plants may justify their wide usage in traditional medicine.

Keywords: Phytochemical analysis, Antimicrobial activity, Medicinal plants, Agar well diffusion method, Leaf extract, Zone of inhibition.

1. INTRODUCTION

Since before recorded history, plants have been used as traditional medicine and remain until now as one of the important resources to combat serious diseases in the world. The discovery of medicinal plants typically resulted from the populace's experience of long and dangerous self-experiments. The better understanding of a plant-derived medicine has depended on two factors [1]. First, the development of increasing strict criteria of evidence that a medicine really does what it is claimed to do and secondly, the identification by chemical analysis of the active compound in plant. About 80% individuals from developing countries rely on plant-based preparations used in their traditional medicinal system and as the basic needs for human primary health care [2]. Plants used for traditional medicine contain a wide range of bioactive molecules, making them rich sources of different types of medicine particularly antimicrobial properties [3]. Mostly, these compounds are secondary metabolites such as alkaloids, flavonoids, steroids, tannins and phenol compounds.

Nowadays, the major problem faced by the world is the multiple drug resistance pathogens (MDR) because of the indiscriminate use of commercial antimicrobial drugs. The massive use of antibiotics in human therapy resulted to bacteria developing several resistance mechanisms since these microorganisms possess the genetic ability to acquire and transmit resistance to therapeutic agents [4]. This resistance problem poses a demand to develop alternative antimicrobial drugs for the treatment of infectious diseases. One approach is to screen local medicinal plants for their potential antimicrobial properties. Antimicrobials of plant origin have enormous therapeutic potential [5]. They are effective in the treatment of infectious diseases while simultaneously minimizing many of the side effects that are often associated with synthetic antimicrobials or antibiotics. There are many published reports on the effectiveness of traditional plants against Gram-positive and Gram-negative microorganisms, and as a result plants are still recognized as the basis for modern medicine to treat infectious diseases [6]. In the present study, five commonly used Surigaonon

medicinal plants *Curcuma longa* (Duyao), *Cymbopogon citratus* (Tangyad), *Coleus aromaticus* (Garabo), *Mentha arvensis* (Helba buena) and *Capsicum frutescens* (Sili) were assessed for their phytochemical contents and antimicrobial activity of their leaf extracts against *Staphylococcus aureus* (gram positive) and *Escherichia coli* (gram negative) bacteria.

2. MATERIALS AND METHODS

Collection of plant materials

The fresh and healthy leaves of the selected medicinal plants used for the experiment were *Curcuma longa* (Duyao), *Cymbopogon citratus* (Tangyad), *Coleus aromaticus* (Garabo), *Mentha arvensis* (Helba buena) and *Capsicum frutescens* (Sili). The plant materials were collected from July to August 2015 from various areas in Surigao del Sur, Philippines. The plant specimens were identified in Biology Department of Mindanao University of Science and Technology.

Preparation of extracts

The plant materials collected were washed thoroughly and dried in shade and powdered in a mechanical grinder. One hundred grams of each respective plant parts were soaked to 300 mL ethanol and kept at room temperature for 24 hours. The extracts were filtered using Whitman filter paper (No.1) and dried at below 45°C for the removal of ethanol for obtaining the dense extract or determining the concentration in mg/mL. The extract was preserved at 2-4 °C and for further investigation for potential antimicrobial properties.

Phytochemical analysis

The extracts of each medicinal plant were subjected to phytochemical testing to ascertain the presence of metabolites such as carbohydrates, reducing sugars, saponins, flavonoids, alkaloids and glycosides by following the standard procedures.

Test for Carbohydrate

The plant extract was mixed with few drops of Benedict's reagent and boiled in water bath. The formation of reddish brown precipitate showed the positive result for the presence of carbohydrate [7].

Test for Reducing Sugar

A 0.5 mL of plant extracts was added with 1 mL of distilled water and 5-8 drops of Fehling's solution and was heated over water bath. Formation of brick red precipitate indicates the presence of reducing sugars [8].

Test for Saponins

The plant extract was mixed with water. It was shaken and observed for the formation of froth, which is stable for 15 minutes. This showed a positive result [7].

Test for Flavonoid

The 4 mg of plant extract was treated with 1.5 mL of 50% methanol. The mixture was warmed and was added with magnesium ribbon. To the mixture, 5-6 drops of concentrated hydrochloric acid was added. Formation of red or orange colour indicates the presence of flavonoids [9].

Test for Alkaloid

The ethanolic extract of plant was evaporated to dryness. The residue was then heated on a boiling water bath with 2% hydrochloric acid. It was then cooled and the mixture was filtered and treated with a few drops of Meyer's reagent. The presence of turbidity or yellow precipitate showed the presence of alkaloid [6].

Test for Glycoside

The plant extract was treated with few drops of glacial acetic acid and few drops of ferric chloride and mixed. The concentrated sulphuric acid was added to the mixture and observed for the formation of two layers. The upper acetic acid layer with a bluish green colour indicated a positive test for glycosides and the lower layer with a reddish brown colour [10].

Isolation of pathogens

Two test pathogens *Escherichia coli* and *Staphylococcus aureus* used in this study were isolated from water samples by standard methods. Identity of the isolates was confirmed by morphological characteristics and conventional biochemical tests [11]. Pure cultures were preserved at 4°C on nutrient Agar.

Determination of antibacterial activity

Antimicrobial activity of five medicinal plant extracts was determined by agar well diffusion method. A 0.1 mL of freshly grown culture of test organisms (106 cfu/mL) was aseptically introduced and spread on surface of sterile Muller Hilton agar plates. Wells of 6 mm diameter were made in agar plate with the help of sterile cork-borer. Fifty microliters of different plant extracts were filled in the wells with the help of micro pipette. Standard reference antibiotics like Chloramphenicol was used as positive controls for the test organisms. Plates were left for some time at 4°C till the extract diffuses in the medium with the lid closed and incubated at 37°C for 24 hr. The plates were observed for zone of inhibition. Antibacterial activity was evaluated by measuring the diameter of the zone of inhibition against the tested bacterial pathogens. Each assay in this experiment was replicated three times [12].

3. RESULTS AND DISCUSSION

In this study, gram positive bacterial pathogen *S. aureus* and gram negative pathogen *E. coli* were selected and antimicrobial activity of leaf extracts of plants *Curcuma longa* (Duyao), *Cymbopogon citratus* (Tangyad), *Coleus aromaticus* (Garabo), *Mentha arvensis* (Helba buena) and *Capsicum frutescens* (Sili) were evaluated against them. The phytochemical screening of the five medicinal plants revealed the presence of various secondary metabolites like carbohydrates, reducing sugar, saponins, flavonoids, steroids, alkaloids and glycosides some of which have been previously associated with antimicrobial activity [13]. These bioactive metabolites are known to act by different mechanism. Like alkaloids, they have shown to possess both antibacterial and antidiabetic properties [14]. Glycosides serve as defense mechanisms against predation by many microorganisms, insects and herbivores [15]. Steroids have been reported to have antibacterial properties [16] and saponin has antimicrobial property which is due to its ability to cause leakage of proteins and certain enzymes from the cell [17].

Table 1: Phytochemical analysis of Ethanol extracts of Selected Medicinal Plants

Secondary Metabolites	Medicinal Plants				
	<i>C. longa</i> (duyao)	<i>C. citratus</i> (tangyad)	<i>C. aromaticus</i> (garabo)	<i>M. arvensis</i> (helba buena)	<i>C. frutescens</i> (sili)
Carbohydrates	+	+	-	+	+
Reducing sugar	+	+	+	+	+
Tannins	-	-	-	-	-
Saponins	+	+	-	+	+
Flavonoids	-	-	+	+	+
Steroids	-	+	+	+	+
Alkaloids	+	+	+	+	+
Anthraquinones	-	-	-	-	-
Glycosides	-	+	+	+	+

(-) absence of secondary metabolites, (+) presence of secondary metabolites done in triplicates

Table 1 revealed that out of 9 metabolites tested *M. arvensis* and *C. frutescens* had the highest number of metabolites followed by *C.citratus* and the least is *C. longa*. The presence of these metabolites is an indicator that the plants can be a potential source of precursors in the development of synthetic drugs

Table 2 revealed that ethanol extracts of *M. arvensis* has highest inhibitory activity against all two test organisms *E. coli* and *S. aureus*. The maximum zone of inhibition (15.537 mm) against *S. aureus* was obtained in *M. arvensis* and the maximum zone of inhibition (15.253 mm) against *E. coli* was obtained in *C. aromaticus*. The present results confirmed the study of [18] that the ethanol extract of *M. arvensis* inhibited the growth of *E. coli* and *S. aureus* including *P. aeruginosa*, *S. flexineri*, *K. pneumonia* pathogens. In addition [19] found in their study that *C. aromaticus* showed more inhibition zone in *E.coli*. *C. aromaticus* and *C. frutescens* showed moderate antimicrobial activity while

C. citratus and *C. longa* extracts showed considerable antimicrobial activity against *S. aureus* (gram positive). For gram negative pathogen *E. coli*, *C. citratus*, *M. arvensis* and *C. frutescens* showed moderate antimicrobial activity and *C. longa* showed less significant in antimicrobial activity. The graph in Figure1 shows the average diameter of zone inhibition of five medicinal plants against both gram-negative and gram-positive bacteria. Results revealed also that the ranged of inhibition of the five medicinal plant extracts are lower than the range of of inhibition of Chloramphenicol as standard. [20] stated that the demonstration of antimicrobial activity against both gram positive and gram negative bacteria by the plants may be indicative of the presence of broad spectrum of antibiotic compounds. Furthermore, the optimal effectiveness of a medicinal plant may not be due to the one the combined action of different compounds originally in the plant [21].

Table 2: Antimicrobial property of Selected Medicinal Plants against Gram positive and gram negative bacterial species tested by disc diffusion assay Zone of inhibition (mm).

Medicinal Plants	Bacterial Pathogens	
	<i>Staphylococcus aureus</i> (Gram positive) mm	<i>Escherichia coli</i> (Gram negative) mm
<i>Curcuma longa</i> (duyao)	7.347±0.560	7.470±0.104
<i>Cymbopogon citratus</i> (tangyad)	7.633±0.064	14.033±0.115
<i>Coleus aromaticus</i> (garabo)	15.000±0.100	15.253±0.040
<i>Mentha arvensis</i> (helba buena)	15.537±0.058	13.627±0.248
<i>Capsicum frutescens</i> (sili)	14.970±0.061	11.117±0.196
Chloramphenicol (500 mg)	23.013±0.110	23.960±0.242

Means of zone inhibition ± standard deviation.




Figure.1 Average diameter of zone inhibition exhibited by the Medicinal Plants against *E.coli* (Gram-negative) and *S. aureus* (Gram-positive).

In addition, the antimicrobial activity of medicinal plants [22] may be due to the presence of different chemical agents which were classified as bioactive antimicrobial compounds. The results of the present study confirmed the validity of the use of these medicinal plants as traditional medicines and also suggests that some of the plant extracts possess compounds with antimicrobial properties that can be used as antimicrobial agents. Use of herbal medicine is safer compared to synthetic drugs. The study scientifically proves the importance of plant products in development of a potent antibacterial agent. However, further studies must be carried out to better evaluate the potential effectiveness of the crude extracts as the antimicrobial agents.

4. CONCLUSIONS

The result of the present study showed that the extract of *Mentha arvensis* which contain the highest number of metabolites, exhibited highest inhibitory activity against two test organisms *E. coli* (gram negative) and *S. aureus* (gram positive). The high antimicrobial activity of *Mentha arvensis* may be due to the presence of secondary metabolites present in plants that provide the necessary component as antimicrobial agent. All of the extracts in this research exhibited different extent of antimicrobial activity.

5. REFERENCE

- [1] Holiman, A., 1989. Plants in Medicine. Chelsea Physic Garden. The Chelsea Physic Garden Co. Ltd.
- [2] Ellof, J.N. 1998. Which extractant should be used for the screening and isolation of antimicrobial components from plants? J. Ethnopharmacol., 60: 1-6.
- [3] Nair, R., Kalariya, T., Chanda, S. 2005. Antibacterial activity of some selected Indian medicinal flora. Turk. J. Biol., 29: 41-47.
- [4] Yala D, Merad AS, Mohamedi D, Ouar Korich MN. Classification et mode d'action des antibiotiques. Médecine du Maghreb, 2001; 91:5-12.
- [5] Cunha, B.A. 2001. Antibiotics side effects. Med. Clin. North Am., 85: 149-185.
- [6] Evans, W.C., 2002. Trease and Evan's Pharmacognosy. 5th ed., Haarcourt Brace and Company, pp 336.
- [7] Raman N. Phytochemical Technique. 2006; New Indian Publishing Agencies: New Delhi p.19. *mexicana* L. (Papaveraceae). CMU J. Nat. Sci., 8: 77-84.
- [8] Iyengar, M.A. 1995. Study of drugs, 8th edn, Manipal Power Press, Manipal, India.
- [9] Siddiqui, A.A., Ali, M. 1997. Practical Pharmaceutical Chemistry. 1st Ed., CBS Publisher and Distributors, New Delhi, pp 126-131.
- [10] Harborne JB. Phytochemical Methods. New Delhi: Springer (India) Pvt.Ltd; 2005. p.17
- [11] Harley, J.P., Prescott, L.M. 2002. Laboratory exercises in microbiology. 5th edn, McGraw Hill, New York.
- [12] Jain, P., Bansal, D., Bhasin, P. 2009. Antibacterial activity of aqueous plant extracts against *Escherichia coli* and *Bacillus subtilis*. Drug Invent. Today, 2(4): 220-222.
- [13] Nweze, E.T., J.I. Okafor and O. Njoku, 2004. Antimicrobial Activities of Methanolic extract of *Trumeguineesis* (Schumm and Thorn) and *Morinda lucinda* Benth used in Nigerian Herb. Medicinal Practice. J. Bio. Res. Biotechnol., 2(1): 34-46.
- [14] Adeshina G.O., Kunle O.F., Onalapo J.O., Ehinmidu J.O., Odama L.E. 2012. Phytochemical and Antibacterial Studies of the Hexane Extract of *Alchornea* Leaf. Phytochemicals as Nutraceuticals – Global Approaches to Their Role in Nutrition and Health. pp. 90-96
- [15] Dhar, M.L., Dhar, M.M., Dhawan, B.N., Ray, C. 1979. Screening of Indian plants for biological activity. Ind. J. Biology., 6: 232-234.
- [16] Raquel, F. Epan, 2007. Bacterial lipid composition and the antimicrobial efficacy of cationic steroid compounds. *Biochimica et Biophysica Acta*, Pp. 2500-2509.
- [17] Zablotowicz, R.M., Hoagland, R.E., Wagner, S.C. 1996. Effect of saponins on the growth and activity of rhizosphere bacteria. *Adv. Exp. Med. Biol.*, 405: 83-95.
- [18] Sugandhi, Rachel Madhuri and Bai Meera. 2011. Antimicrobial Activity of *Mentha Arvensis* L. *Lamiaceae*. *Journal of Advanced Laboratory Research in Biology*.
- [19] Subhas, Chandrappa M, Harsha, Dinesha and Thammanna, Gowda. 2010. Antibacterial Activity of *Coleus Aromaticus* Leaves. *International Journal of Pharmacy and Pharmaceutical Sciences*, ISSN- 0975-1491 Vol 2, Issue 3.
- [20] Evans CE, Banso A, Samuel OA; *J Ethno pharmacology*, 80 2002, pp. 21–24
- [21] Joshi, B., Sah, G.P., Basnet, B.B., Bhatt, M., Sharma, D., Subedi, K., Pandey, J., Malla, R. 2011. Phytochemical extraction and antimicrobial properties of different medicinal plants: *Ocimum sanctum* (Tulsi), *Eugenia caryophyllata* (Clove), *Achyranthes bidentata* (Datiwan) and *Azadirachta indica* (Neem). *J. Microbiol. Antimicrob.*, 3(1)
- [22] Arulmozhi, S., Mazumder, P.M., Ashok, P., Narayanan, L.S. 2007. Pharmacological activities of *Alstonia scholaris* Linn. (Apocynaceae)- A review. *Pharmaco. Rev.*, 1: 163-165.